

The serotonin model of suicidal propensities¹

Michael Heller

In all civilisations of the world there are people who kill themselves. It would seem that the social systems which generate the greatest number of suicides are the industrial societies generated by Europe (capitalist or communist). Only Sri Lanka and Surinam present suicide rates as high as those observed in Europe. Switzerland does not escape that trend, as it was, in 1987, classed 9th by the World Health Association in terms of world suicide rate. Furthermore in our societies the suicide rate tends to increase dramatically since 1965. During the 19th century (Durkheim 1930), suicide rate increased with age. The new increase corresponds quasi exclusively in an increase of suicides among males from 15 to 24 years of age. Suicide attempts yield much higher numbers. In 1976 427 persons out of 100 000 (431 174 cases) in the EEC made a suicide attempt. (Diekstra 1989)

In 1971 the Geneva statistics mention 58 suicides in the year (Service cantonal de statistique de Genève 1971). Today, in the same town, the number is approximately a 100 a year. As a culture we are clearly addressed by such figures, and it is our responsibility to study the factors — clearly social or psychological via education — which support more and more often the citizen's death wishes. It is also well known that these official figures are probably inferior to reality, as an unknown number of accidents are probably masked suicides, or suicide attempts.

In our culture it is usually medical institutions who are asked to judge the state of persons who attempt suicide. Furthermore it has often been observed that many suicides consulted for medical help a few months before their death (Yessler and all. 1961, Brown & Sheran 1972, Nielsen, B., Wang, A.G. & Bille-Brahe, U. 1990). In

¹ This is a draft version of the introduction I wrote for a "Request made to the Suisse National Funds for Research in Psychiatry, made in 1991 for the suicide study of Marc Archinard, Michel Heller et André Haynal". The project was accepted as n.3233546.92. It supported a research carried out by the *Laboratoire Affect et Communication* (L.A.C.) created by André Haynal and myself in the Geneva University Psychiatric Institutions (I.U.P.G.) and the Geneva Department of medicine in 1988. The results are summarized in Heller, Michael; Haynal-Reymond, Véronique; Haynal, André & Archinard, Marc (2001): *Can Faces Reveal Suicide Attempt Risks?* Michael Heller [ed.], *The flesh of the soul: the body we work with*, pp. 231-256. Bern: Peter Lang. I do not have the final text. A few corrections were added on October 24 2006.

This publication on internet is made available for private use only. Any form of organized distribution of these texts requires the permission of the authors, as expected by the copyrights.

the more recent structures it is observed that from 30 to 64 percent of suicides had a recent contact with psychiatric institutions (Wolk-Wasserman, D. 1987). Niels, Wang & Bille-Brahe and (1990) report that in the Denmark of 1987 psychiatric institutions have seen 10 000 suicidal patients.

Given this state of affairs many studies have attempted not only to understand the mechanisms that lead to suicide, but also methods of finding the suicide risk for an individual. All models and methods found until today, although helpful, are notoriously insufficient. Recently, having mentioned a general agreement "that the estimation of (suicide) risk is important, complex, and difficult to quantify." Motto (1991) summarizes the present situation by stressing that "the absence of generally accepted instruments to accompany the task attests to the elusiveness of a unitary solution to the problem."

Ways of understanding suicide

Suicide has been approached through many angles, ranging from sociological models to physiological ones. The literature can be summarized as oscillating between two extreme positions:

- a) A literature which stresses the variety of causes, motives, and type of persons who commit suicide. Starting with Durkheim's (1930) classical study, this literature shows that suicide rates are influenced by social factors such as religion, geography, and marital status independently from psychopathology.
- b) Although convincing, these data are of little help for the clinician. It has indeed been confirmed that a northern protestant bachelor living in town has more chances than a Mediterranean catholic married in a village to suicide. But the figures we are speaking of are too small for the clinician, for it is obvious that many northern protestant bachelor living in town do not suicide. Running away from the sociologist's defence of diversity, some clinicians hoped that a form of suicidal predisposition could be detected in individuals. This has led to concepts of a 'suicidal syndrome' (Ringel 1976), or 'suicidal profile' (Nielsen, B., Wang, A.G. & Bille-Brahe, U. 1990, Motto, J.A 1991).

Current research seems to move between these two polarities. For example Freud's proposition that depression may be a major factor in suicide not only strengthened intra-psychic explanations, but influenced sociological research to focus on depressing social

factors such as delinquency, broken home, unemployment, place of residence, unemployment, job loss, school and professional failure (Brown & Sheran 1972, Rourke, Young, Leenars 1989)... which is not what sociologists had in mind. This cognitive loop, induced by the preoccupation of medical institutions, has a certain wisdom, as it reflects problems indeed encountered by a probably important part of the suicidal population which consult for medical support: people who have suicidal impulses, who are not able to contain them, and some times act upon them without having really made the choice of dying. These people are the main preoccupation of clinicians, as it is in a certain sense only in those cases that medical support is really required and ethically essential. But one must not forget many other cases which probably addresses moral and/or ethical issues more than purely medical ones.

The prevalent medical position relating suicidal intentions to psychopathology does not really take into account people hoping to avoid being transformed in pluming factories by hospitals, people attempting to avoid speaking under torture, or people killing themselves under the influence of social rituals such as hara-kiri in Japan. But even opposing 'moral' suicides to those related to psycho-pathology is arguable, as:

- a) some people do not speak under torture
- b) people with a clear psychopathology seldom suicide.

The serotonin hypothesis

An important problem for psychiatry, often stressed by justice institutions and medias, is that of predicting who will commit and overt aggressive act against others or self under what appears to be a 'mad impulse'. People just released from a psychiatric institution who attack a whole village, or the children of region are particularly dramatic examples. Suicides are more current, although less overtly embarrassing, cases.

Today this problem is approached around a central theme: the relations between serotonin and impulsivity. Several studies have shown serotonin levels to be lower among people with impulsive (auto) aggressive acts than among current depressives, who themselves have a lower serotonin level than a 'normal' population. In the last five years research has provided an important number of studies confirming this study (Iraskman-Bendz L., Asberg M., Nordstrom P., Stanley M. 1989, Montgomery S.-A. 1987, Moss H.B., Yao J.K. & Panzak G.L. 1990, Ricci L.-C., Wellman M.-M. 1990), and one study in which this finding was not replicated (Roy

A., Lamparski D., de Jong J.-A., Adinoff B. & al. 1990)². Although a serotonin explanation of impulsivity will probably soon prove to simplistic, it is for the moment the hypothesis which groups a maximum of data.

A first way of dealing with the serotonin findings is to consider them as a blunt correlation, which would require other correlations to be related to more general mechanisms. We prefer to spell why the serotonin model is appealing as a finding fitting in a whole other series of other experimental and clinical findings, which for the moment have not been connected with the serotonin findings empirically. And then to consider how those relations could be tested experimentally.

The ethological perspective

Social status in schools

Difference of social status, in an ethological perspective, is considered as an asymmetric relation that can be organized in a hierarchy in function of dominance. (...) Generally dominance is defined in function of three dimensions: (i) access to resources and sexual partners, (ii) who wins fights and (iii) behaviors stating the others' status (submissive behavior, who looks at who, presentation of genitals, etc...). In fact a given primate can not always be classified in the same way in relation to these three dimensions: a person can be very dominant in relation to scale (i), and much less dominant in relation to scale (iii), etc... (Feyereisen & De Lannoy 1985: 33-34, 210-211).

Ethologists have extended their approach of dominance to the study of human groups such as school classes. From this perspective status display has been related to ways of asserting "territorial rights, such as putting a foot or leg on a desk drawer, desk top or chair, touching or leaning against property, distributing personal articles to mark territory, crowding what is ostensibly someone else's space or property as in another's office" (Henley 1977:127). This approach of leadership has been applied by Montagner (1978) in a study where he analyzed in detail the behaviour displays of school-boys, and found the following distinction:

Les leaders expriment des actes de lien et d'apaisement beaucoup plus souvent que les dominants-agressifs. Les leaders provoquent ainsi chez les autres enfants l'approche et l'offrande, sans qu'il soit possible de conclure qu'ils ont exprimé de telles attitudes dans l'intention d'attirer et d'entraîner les autres. L'enfant de 2-3 ans est d'autant plus attirant, imité et suivi que la fréquence de

². On the other hand relations between suicide and catécholamines or the dexamethasone suppression test does not seem to be confirmed.

ses séquences d'apaisement est plus élevée et celle de ses agressions plus faible (p.161).

The attitudes used to describe a status are defined as corresponding to established bodily patterns (listed in pages 34 — 37) such as caress, kiss, touching another's hand, to put one's hand on another's neck, to lean one's head laterally... (p.90-94).

Nadel-Brulfert, Baudonnière, & Fontaine (1983) show that Montagner's patterns are only observed when many children are confronted with a small number of toys. In situations where three children are put in a room with three samples of each toy, 'monkey dominance behaviors' are not observed. This experiment suggests that such dominance behaviours as those observed by ethologists among humans, can often be related to situational phenomena (socio-psychological), rather than more sociological ones.

The Nadel-Brulfert study suggests that 'apish' dominance behaviours tend to intrude in the dynamics of a group when its members have to fight to find what they need.

This literature constantly distinguishes a 'leader' and an 'impulsive' character. The impulsive character is often aggressive in an inefficient way, which associates him with what family therapists call a 'symptom child', and which psychoanalytical literature associates with a 'weak ego' structures (psychotic, border-line, narcissic...). These two types of character can be highly aggressive, but they do not manage their aggressivity in the same way:

— The leader character is described as some one who prefers to display appeasing gestures, and only displays aggressivity when it is required by both personal and situational needs. When these needs are frustrated, the leader character finds efficient ways of destroying his enemies. (Laborit 1979, p.116)

— The impulsive character, on the other hand, is described as having a less appeasing behaviour, and as displaying highly aggressive gestures very quickly, even when it does not seem to be required for his personal or situational benefits. Further more he is described as some one who is less strong than a leader, and who tends to loose fights against him. (Montagner 1978)

It can therefore be postulated that the leader has a good capacity to master and regulate his aggressivity, while the 'weak aggressive' has a bad capacity of regulating it. Leadership, in ethological literature, mostly considers a form of natural leadership character, which does not have to include such human phenomena as Napoleon or Hitler.

Serotonin and dominance

Buck (1988, p.213) describes an experiment of McGuire, Raleigh, and Brammer demonstrating "an interesting similarity in the levels of serotonin in dominant male monkeys and dominant male humans. They found that the level of serotonin in the blood of

dominant male rhesus monkeys is twice as high as the level in other male members of the group. When the dominant male is removed from the group, his level of serotonin falls to normal, whereas the level of serotonin in the newly dominant male rises. According to the evidence, the rise in the leader's serotonin level is induced by the submissive behaviour he receives from the followers. A similar phenomenon was found in the blood serotonin levels of members of a fraternity at the University of California at Los Angeles: the fraternity officers had higher levels than other members. Whether shifts in leadership status affect the serotonin levels of male humans remains to be investigated. It is noteworthy that McGuire and his colleagues found dominance to be associated with *high* levels of serotonin, for impulsive violence seems to be associated with *low* levels of the same substance. The crucial difference here may be between confident dominance and 'bad-tempered' hostility. "

If one considers the initial functions of serotonin, this result is indeed astonishing one. Serotonin was initially perceived as an inhibitor of behaviour, "inhibitory and restorative of general bodily resources. Serotonin is related with the control of the parasympathetic nervous system." Inhibition of serotonin causes insomnia, whereas increased amounts of serotonin appear to increase slow wave sleep. Decrease of serotonin correlates with increase of motivated behaviour; increase of serotonin may be related to increase of introversion. (Buck 1988, p. 137, 139, 145ss, 549ss)

Laborit (1990, p. 118ss) points out that the serotonin effect must be perceived as balancing others psycho-biological effects if one wants to understand it, and shows how complex a system present knowledge already points to. For example (p.123), among monkeys, leaders tend to use the adrenal medullar system when involved in social interactions, while subordinate males who are less active tend to respond with the hypothalamus-pituitary-adrenal cortex axis... known to be involved in stress responses. Serotonin has the effect of diminishing action. As such it can counterbalance a propension to move, activated by other systems such as those influenced by catecholamines. Leaders from this perspective, are described as (a) having a high charge, and (b) as having the means (enough serotonin is available) to put strong breaks on that charge.

I have not found literature showing that leader and impulsive characters have a different aggressive potential. What mainly differentiates them is their capacity to inhibit impulsive aggressive behaviour. This is stressed by the observations that changing certain biological variables allows different monkeys to become leader, as shown not only in this experiment, but already in the classical experiment of Plotnik, Mir, and Delgado (1971), were

monkeys could themselves stimulate electrodes planted in certain nervous centres.

Another aspect of the above mentioned research, is that serotonin level and ability for leadership is regulated — among monkeys — by submissive behaviour. This presupposes a regulative system between 'leadership behaviour' and 'submissive behaviour'. In a certain way one could say that a leader needs to eat the other's serotonin. This is accomplished by behaving in such a way that others offer submissive behaviour. Thus the serotonin model supposes that one of the variables influencing serotonin concentration in an organism is interaction between behaviour on a very specific axis: dominance — submissive. Becoming the centre of gratifying displays in a population creates in me the bio-psychological possibility of becoming a leader of this population (thus one can become the leader of a family which has adored one, and not a leader in an other population — such as a professional one — were we did not manage to receive gratifying displays. In more general terms dominance capacities in a population is regulated by distribution of gratifying displays; and an incapacity of eliciting such displays would create a negative loop on some one's capacity of dominance. (Buck 1988, p. 213)

The psychoanalytical perspective

Ego strength and impulsivity

In 1917 already, Freud (1953, p. 252) had noticed intense "murderous impulses against others" "turned back" on oneself. But he also found that aggressivity directed to oneself was not a sufficient cause "to explain what interplay of forces can carry such a purpose thought to execution". Sociology had already shown that there is a much bigger number of people who can be classed in suicide risk populations than the number of people who actually suicide. Freud confirms at a psychological level that there are many more people with high auto-aggressive and suicidal impulses than the number of people who carry out a suicidal intention³.

To explain which people among those with suicidal fantasies might finally carry them out, psychoanalytical 'characteriology' also distinguishes between weak and strong ego. One of the functions of the ego, is the regulation of our instinctual and emotional life from at least three points of view:

³. I have not found a study showing that at least most suicides occur in those populations with a high risk.

a) A *delay* system: expressing an emotion or satisfying an instinctual urge must often be delayed until we find our selves in a suitable situation. This delay system must at least allow for regulations within a day.

b) A *cooling* system: the delay system can only function if our emotional psyche can be somewhat cooled for a moment so that for example we can concentrate on our work until lunch time even if an attack of hunger intrudes in our mind at eleven o'clock.

c) A *recycling* process: when an emotion can not be adequately expressed within a matter of days, the organism has the possibility of expressing an urge through less appropriate channels. For example sometimes it may be more appropriate to use express anger by overworking than by shouting at one's boss and losing one's job.

Two types of psychopathologies can be derived from this model:

a) people who only have some of these means available, and therefore over-use them (neurotic pole)

b) people for whom all these means are insufficiently developed (psychotic pole)

When ego is too weak for one or more of these functions people tend to rely on a more voluntary form of control: will power, generating what one often calls 'super-ego' pathologies. The term 'will power' itself shows the limits of such a mode of self regulation: it is strongly dependent of the amount of 'inner power' available at each moment in us. Thus during moments of inner weakness, the 'wilful person' finds himself incapable of regulating his behaviour in a coherent way, and may act upon impulses⁴. Acting on impulses may mean doing things parts of myself wish to do, but which other parts of myself might not want to do. Thus, reacting to a beautiful but depressing film I might kill myself, while more deeply I might wish to live. It is to be noticed that in this model one can find impulsive aggressive acts among most character structures (as all have at least one weak function), but that it becomes more likely as one comes nearer to the psychotic pole. This suicide distribution corresponds to what is generally observed.

Following such a line of thought, Rainer Krause, who helped us to put up our experiments since 1987, had suggested that a likely area to find behavioural traits related to suicide would be in the direction of impulsive acts.

Serotonin and psychopathology

Psychiatric studies have confirmed a relation between quantity of available serotonin, structures considered as having a weak ego by psychoanalysts, and impulsive (auto) aggressive acts. This trend of research tends to favour a relation between suicide and low serotonin rather than a relation between suicide and depression. Edman, Asberg, Levander & Schalling (1986) clearly

⁴. Reich.

specify "the association (of serotonin level) with suicide, however, is not confined to melancholia, as suicide attempters who do not satisfy research criteria for depression have lower CSF 5-HIAAA levels⁽⁵⁾ than control subjects. Furthermore, subjects with personality disorders and borderline conditions who have attempted suicide had lower CSF 5-HIAAA levels than those who had not." Moss, Yao, & Panzak (1990) stress that "alterations in serotonergic neurotransmission have been implicated in disorders characterized by impulsivity, aggression, low mood, and substance abuse." They found low serotonin concentration among antisocial personality disorders (ASP) as well.

The relationship between CSF 5-HIAA and suicidal behavior appears to be most clear cut for suicidal acts, in particular when violent methods have been used. Thus, when suicidal ideation was rated or a rating scale was used that combines ideation and acts, the association was weaker or nonexistent (Edman, Asberg, Levander & Schalling, 1986).

Brown and Goodwin (1986, p. 144-145) stress that low serotonin levels are associated first of all with violent acts, and then with suicide; with impulsiveness and an inability to "delay gratification". It would therefore seem that an impulsive violent act requires (a) a tendency for violence, and (b) a low serotonin. In other words it would appear as if serotonin could be related to the cooling function of the psychoanalyst's ego, and that further research on this model could help psychiatrists to detect people who could end up committing a violent act.

Serotonin findings are probably part of a more general mechanism. For prevention and therapeutic purposes it could be helpful to find other elements of this general mechanism. The literature provides us with clinical observations which a) could fit with the model we already constructed around the serotonin findings, b) are considered helpful in dealing with depressive patients, and c) add new elements to an eventual model of the suicidal population which could benefit from psychotherapeutic and psychiatric help: those for who may commit suicide on the basis of a whim, rather than as a voluntary and responsible decision.

Suicide and significant others

The more sociologically oriented studies (Brown & Sheran 1972) suggest that "suicide is probably best understood as a multidimensional malaise", through which biological, sociocultural, interpersonal, psychological, and personal/existential aspects are coordinated (Rouke, Young, & Leenars 1989). It has been noticed that this coordination of levels involves at least:

⁵. 5-hydroxyndoleactic (a substance close to serotonin) acid concentration in the cerebrospinal fluid.

- a) Depressing situations such as unemployment, job loss, school and professional failure...
- b) inability to cope or solve problems in general
- c) marked interpersonal conflict and especially a feeling of rejection.

These observations already set a scene which is not gratifying for one's self-esteem in terms of dominance. Observations on the relations between suicidal patients and the 'significant others' they live tends to increase the feeling of persons were feed-back can be perceived as negative.

Fawcett, Leff & Bunney 1969 quote clinical observations showing that "suicidal behavior occurs in a symbiotic relationship of mutual dependence and masochism". Wolk-Wasserman (1986) observed that family members manifested extremely strong hostility and expressed death wishes against the patient. He suggests that suicide occur with people who have to deal with strong displays of aggression around them. Wolk-Wasserman (1987) also summarizes observations by Richman who "stated that the suicidal family is characterized by its closed nature and broken links with social institutions, as with friends and acquaintances. Family members were therefore reduced to seeking help within the closed family circle. If this support failed, there was a risk of suicidal action".

A suicidal person would then be (a) over-dependent of a 'tribe' which provides an aggressive and dismal feed-back, and (b) a person who is aggressive against herself and has a dismal view of what can be lived in this world. Among those we should find most suicides, as such a setting would tend to lower serotonin levels.

Suicidal syndrome

Ringel (1976), influenced by psychoanalysis, attempts to describe a sort of 'suicidal character'. The core feature would be what Ringel calls dynamic constriction. Constriction is an impression of being constricted. Dynamic constriction also has the following properties:

A compulsive urge into one direction, that is, in the direction of ultimate suicide. (...) In the outward appearance of the individual, dynamic constriction is frequently manifested by a loss of spontaneity, by inhibition and passive behavior. (...)The patient feels that the situation or the condition will never change and thus he falls prey to his own mood.

There is a general depreciation of self and others. Ringel also observes a strong super-ego pathology, full of aggression against the self. He also stresses that suicide is independent from specific pathologies.

Motto (1991) finishes this portrait by noticing the difficulty of such people with pain: "Suicide occurs when a person's limit of psychic pain tolerance is exceeded by the level of pain that is either experienced or anticipated with a degree of subjective certainty. Thus the process of estimating the risk of suicide at a given

moment is essentially determining how close the actual or anticipated pain is to the person's tolerance threshold at that time."

Discussion

Testing serotonin levels is today a very tedious task which is difficult to carry out on such people as suicidal patients who come for help and support after an attempt. Nevertheless these studies have helped us to group a certain number of isolated observations, and propose the skeleton of a model we wish to test in the future.

What we have tried to convey, is model involving processes going from the biochemical to the social, and from the social to the biochemical. All of the literature seems to agree on this, yet little has been done on this matter in terms of close observation. The importance of going with research based on such premises is more and more often stressed in the literature on emotional life and emotional difficulties (Buck 1988, Haynal 1990).

We do not have the means to carry out a research strategy which would allow us to study the coordination of these levels simultaneously, although we do think that such a strategy would be required.

As a compromise solution we have asked ourselves if there was an aspect of this model which has been particularly left aside. And we noticed that one such aspect is nonverbal communication.

The McGuire experiment, and the systemic approaches, both show the difficulty impulsive people have in establishing an efficient relation with others in terms of their self-esteem. These patients seem to have a behaviour which leads them to a dead-end. Therapeutically this means that by giving serotonin to such patients only, without the support of gratifying relationships would be inefficient; and *visa versa*.

Wolk-Wasserman (1987) has made an observation on the relation between five prepsychotic/psychotic suicidal patients and psychiatric services which goes in the same direction:

On the day preceding the suicide attempt, all five patients conveyed their suicidal intentions by indirect, non-verbal communication. This suicidal communication could be expressed by the patient — although in regular contact with the therapist — suddenly arriving at, or repeatedly ringing, the psychiatric department, driven by a need to see the therapist and make sure the contact would continue. This uncertainty regarding the continued contact, and the suicidal thoughts arising in this connection, were often caused by a misapprehension, or misinterpretation of the therapist's remarks about concluding the contact.

In all the cases when the patients arrived unannounced, they were received by the personnel on duty who, failing to understand the indirect, nonverbal suicidal communication and seriousness of the situation, referred to ordinary consultation hours or regulations in force. ... The therapist, after receiving the messages from the duty personel, like the relatives, understood

the indirect, non-verbal suicidal communication and the seriousness of the situation, were upon thy immediately contacted the patients, and also arranged the hospital care.

From such reports one can only conclude that a detailed explicit knowledge of the non-verbal communicative strategies of patients with suicidal risk could be very helpful.

Bibliography

- Brown G.L. & Goodwin F.K. 1986: Human aggression and suicide. In R. Maris. (Ed.): *Biology of suicide*. New York: The Guildford press.
- Brown T.R. & Sheran T.J. 1972: Suicide prediction: a review. *Life-threatening behavior*, Vol. 2, No 2, summer, p.68-97.
- Diekstra R.F.W. 1989: Suicide and the attempted suicide: an international perspective. *Acta Psychiatr Scand*, vol. 80, p.1-24
- Durkheim E. 1930: *Le suicide*. Paris: Presses Universitaires de France, 1990.
- Edman G., Asberg M., Levander S. & Schalling D. 1986: Skin conductance habituation and cerebrospinal fluid 5-hydroxyindoleacetic acid in suicidal patients. *Arch. Gen. Psychiatry*, vol. 43, June, p.586 — 592.
- Fawcett, J., Leff, M. & Bunney, W. E. 1969: Suicide. *Arch Gen Psychiatry*, Vol. 21, p129-137
- Feyereisen P. & De Lannoy J-D. 1985: *Psychologie du geste*. Bruxelles: Pierre Mardaga.
- Freud S. 1953: Mourning and melancholia. In J. Strachey (Ed.): *Standard edition of the complete psychological works of Sigmund Freud*, vol 14. London: Hogarth press.
- Ganong W.F. 1991: *Review of medical physiology. Fifteenth edition*. Englewood cliffs: Prentice Hall.
- Gibbs A. 1990: Aspects of communication with people who have attempted suicide. *Journal of advanced nursing*. volume 15, n 11, p.1245-1249.
- Hassanyeh F., O'Brien G., Holton A.R., Hurren K., Watt L. 1989: Repeat self-harm: an 18-month follow-up. *Acta Psychiatr Scand*, vol. 79, p.265-267.
- Haynal A. 1990: *Psychanalyse et science face à face*. Mezieu: Césura Lyon Edition.
- Heller M. 1992: *Social status and postural dynamics*. In print.
- Henley N.M. 1977: *Body politics. Power, sex and nonverbal communication*. Englewood Cliffs: Prentice-Hall.
- Iraskman-Bendz L., Asberg M., Nordstrom P., Stanley M. 1989: Biochemical aspects of suicidal behavior. *Progress in neuro-psychopharmacology and biological psychiatry*. vol. 13, p. 535-544.

- Kovacs, M., Beck, A.T. & Weissman, A. 1976: The communication of Suicidal Intent. *Arch Gen Psychiatry*, Vol. 33, p.198-201
- Laborit H. 1979: *L'inhibition de l'action. Biologie, physiologie, psychologie, sociologie*. Paris: Masson.
- Laborit H. 1990: *Les récepteurs centraux et la transduction des signaux*. Paris: Masson.
- Lecrubier Y. 1988: Sérotonine et capacité à différer une conduite: un modèle psychobiologique. *Psychiatrie & psychobiologie*, vol. 3, may.
- MacNaughton N. 1989: *Biology and emotion*. Cambridge: Cambridge university press.
- Montagner H. 1978: *L'enfant et la communication*. Paris: Stock.
- Montgomery S.-A. 1987: The psychopharmacology of borderline personality disorders. *Acta-psychiatrica-belgica*, vol 87, n.1, p.260-266, may-june.
- Moss H.B., Yao J.K. & Panzak G.L. 1990 Serotonergic responsivity and behavioral dimensions in antisocial personality disorder with substance abuse. *Bio. psychiatry*, vol. 28, p.325-338.
- Motto, J.A 1991: An integrated approach to estimating suicide risk. *Suicide-Life-Threat-Behav.* 1991 Spring, Vol. 21 (1), p.74-89
- Murphy G.E. 1984: The prediction of suicide: why is it so difficult ? *American journal of psychotherapy*, vol. XXXVIII, No 3, July, p.341-349.
- Nielsen, B., Wang, A.G. & Bille-Brahe, U. 1990: Attempted suicide in Denmark. (IV. A five-year follow-up). *Acta Psychiatr Scand*, vol. 81, p.250-254
- Plotnik R., Mir D. & Delgado J.M.R. 1971: Aggression, noxiousness and brain stimulation in unrestrained rhesus monkey. In Eleftherion B.E. & Scott J.P. (Ed.): *The physiology of aggression and defeat*. London: Plenum Press.
- Ringel E. 1976: The presuicidal syndrome. *Suicide and life threatening behaviour*, 6, p. 131-149.
- Rourke B.P., Young G.C. & Leenars A. A. 1989a: A childhood learning disability that predisposes those afflicted to adolescent and adult depression and suicide risk. *Journal of learning disabilities*, vol. 22, n. 3, p.137-200, March.
- Roy A., Lamparski D., de Jong J.-A., Adinoff B. & al. 1990: Cerebrospinal fluid monoamine metabolites in alcoholic patients who attempt suicide. *Acta Psychiatrica Scandinavia*, vol 81, n.1, january, p.58 — 61.
- Schmidtke, A., Fleckenstein, P. & Beckmann, H.1989: The dexamethasone suppression test and suicide attempts. *Acta psychiatr scand*, vol. 79, p.276-282
- Ricci L.-C., Wellman M.-M. 1990: Monoamines: biochemical markers of suicide? *Journal of clinical psychology*, vol. 46, p. 106-116, january.

- Service cantonal de statistique de Genève 1971: *annuaire statistique 1971*. Genève: Département du commerce, de l'industrie et du travail.
- Thorell L.-H. Kjellman B.F. & Elia G. 1987: Electrodermal activity in antidepressant medicated and unmedicated depressive patients and in matched healthy subjects. *Acta psychiatr scand*, vol. 76, p.684-692
- Thorell, L.-H. 1987: Electrodermal activity in suicidal and nonsuicidal depressive patients and in matched healthy subjects. *Acta psychiatr scand*, vol. 76, p.420-430
- Wolk-Wasserman, D. 1986: Suicidal communication of persons attempting suicide and responses of significant others. *Acta Psychiatr Sand*, vol.73, p.481-499
- Wolk-Wasserman, D. 1987: Contacts of suicidal neurotic and prepsychotic/psychotic patients and their significant others with public care institutions before the suicide attempt. *Acta Psychiatr Sand*, vol. 75, p.358-372
- Yessler, P. G., Gibbs, J. J. & Becker, H. A. 1960: On the Communication of Suicidal Ideas (I. Some Sociological and Behavioral Considerations). *Arch Gen Psychiatry*, Vol. 3, p.60-79
- Yessler, P. G., Gibbs, J. J. & Becker, H. A. 1961: On the Communication of Suicidal Ideas (II. Some medical Considerations). *Arch Gen Psychiatry*, Vol. 5, p.34-51